

$$F_{\nabla} = 2\pi \cdot r^3 \frac{\sqrt{\epsilon_B}}{c} \left( \frac{\epsilon - \epsilon_B}{\epsilon + 2\epsilon_B} \right) (\nabla \cdot I)$$

$F_{\nabla}$  = Optical force on particle towards higher intensity

$r$  = Radius of particle

$\epsilon_B$  = Dielectric constant of background medium

$\epsilon$  = Dielectric constant of particle

$I$  = Light intensity ( $\text{W}/\text{cm}^2$ )

$\nabla$  = Spatial derivative

**FIG. 1**

FIG. 2





FIG. 3






FIG. 4



**FIG. 4A**










FIG. 6





**FIG. 9A**



**FIG. 9B**





FIG. 10








FIG. 12A




FIG. 12B




FIG. 12C




FIG. 13A




FIG. 13B

FIG. 13C




FIG. 14A




FIG. 14B




FIG. 15




FIG. 17




FIG. 16




FIG. 18







FIG. 20





*FIG. 22*




FIG. 23




FIG. 24





FIG. 25

Before:

SCATTER FORCE SEPARATION



After:




FIG. 26

FIG. 27A





FIG. 27B




FIG. 27C





FIG. 28

FIG. 29





HEMOGLOBIN - O<sub>2</sub> ABSORPTION SPECTRUM

FIG. 31






FIG. 32





FIG. 33



Before

After

Difference

FIG. 34

DISTRIBUTION OF ESCAPE VELOCITIES  
READING TAKEN IN PBS/1% BSA BUFFER  
RAIN-X COATED SLIDE/CYTOP COATED COVERSIP





FIG. 35



*FIG. 36*